



INFECTION CONTROL

Introduction

An infection is a disease or illness caused by micro-organisms (infectious agent) such as bacteria, viruses, fungi, and parasites. People with disability are often at a greater risk of infection particularly if they have a chronic disease or are frail. This policy aims to reduce the risk of infection both to clients and to employees.

Infections require three main elements to spread:

- A source of the infectious agent
- A mode of transmission
- A susceptible host

This is the chain of infection and breaking the chain of infection helps to stop the spread of disease. Common modes of transmission include contact, droplet, and airborne. Some disease can spread by more than one mode of transmission, e.g., flu can be spread by droplet or contact.

When supporting people with disability, PPE (personal protective equipment) is often used and depending on the support need can include:

- Gloves
- Gowns or aprons
- Face masks
- Protective eyewear

Applicability

When

- Applies at all times

Who

- Applies to all employees including key management personnel, full time employees, part time employees, casual employees, contractors, and volunteers

Infection prevention

Stopping the spread of infection is everyone's responsibility and includes:

- Practising good hand hygiene
- Following respiratory hygiene/cough etiquette
- Wearing PPE where required
- Correctly handling medical devices
- Correctly cleaning and managing spills
- Correctly handling food, waste, and linen

Hand hygiene

Hand hygiene is the single most important factor in reducing the spread of infections. Hand hygiene can be performed either by washing hands with soap and water or using an alcohol-based hand rub.

The five moments for hand hygiene:

1. Before touching a participant or their surroundings
2. Before a procedure or where there is risk of being exposed to body fluids
3. After a procedure or body fluid exposure risk
4. After touching a participant
5. After touching a participant's surroundings (e.g., bed, linen)

Additionally, hand hygiene should always be performed:

- Before putting on gloves
- After removing gloves
- Before touching food and eating
- After going to the toilet
- After lunch breaks or other breaks
- After blowing your nose or coughing
- After handling waste
- After handling unwashed linen or clothing
- After handling animals
- When your hands are visibly dirty

Respiratory hygiene/cough etiquette

Covering sneezes and coughs prevents infected people from dispersing droplets into the air where they can spread to others. Practicing good respiratory hygiene means:

- Covering your nose and mouth with a tissue when you cough, sneeze, wipe or blow your nose.
- If no tissues are available, cough or sneeze into your elbow rather than your hand.
- Encouraging clients to use tissues when they sneeze or cough.
- Providing the means for prompt disposal of used tissues in general waste.
- Encouraging clients to practise hand hygiene.
- Making alcohol-based hand rub available.

PPE – gloves

Gloves should be worn:

- For procedures with a risk of exposure to blood or body substances i.e., assisting a client with toileting, urinary catheter care or when dressing wounds
- When touching equipment or surfaces that may come into contact with blood or body substances
- When performing blood glucose monitoring
- When performing oral suctioning
- When caring for clients who have an infection spread by contact
- If the employee has broken skin
- When preparing food

Remember:

- Gloves are not used instead of hand hygiene
- Perform hand hygiene before and after using gloves
- Remove gloves when a care activity is finished, change gloves before starting a different care activity
- Dispose of used gloves immediately
- Do not use multiple gloves at the same time.

PPE – face masks

Face masks protect an employee's nose and mouth from exposure to infectious agents and are used if there is a risk of:

- Droplets or aerosols (e.g., from coughs or sneezes)
- Splashes or sprays of blood or body fluids (e.g., when providing complex wound care or urinary catheter care)

Face masks protect the wearer when caring for participant who has an infection spread by droplet or airborne route. Face masks can also be worn by clients who are coughing, especially if they are unable to cover their mouth (both only if wearing the face mask will not cause the client distress).

When using face masks:

- Check the manufacturer's instructions before use.
- Don't touch the front of the mask with your hands once the mask is in place.
- Use the mask for care of one person only and change if a care activity is taking a long period of time.
- Don't leave the mask dangling around your neck.
- When the care activity is complete, discard mask and perform hand hygiene.

PPE – protective eyewear

Protective eyewear protects an employee's eyes from exposure to infectious agents.

when there is a risk of:

- Droplets or aerosols (e.g., from oral suctioning)
- Splashes or sprays of blood or body fluids (e.g., when emptying urinary catheter bags)
- Protective eyewear protects the wearer when caring for participants who have an infection spread by droplet or airborne route.

When using protective eyewear:

- Remember that the outside of the eyewear is contaminated.
- When care is complete, remove eyewear using the headband or earpieces.

- Clean eye shield after each use with detergent and water and allow to dry.
- If eyewear is single use, dispose after completion of care activity.

Handling medical devices

When supporting clients such as with urinary catheters or tracheostomy, good practices are vital to protect both the participant and the employee from risks of infection.

When supporting clients with medical devices:

- Always perform hand hygiene before any contact with the device or where the device enters the clients body.
- Wear PPE if there is risk of exposure to blood or body fluids.
- Touch the device as little as possible.
- Remember that the longer a device is in place, the greater the risk of infection.
- Medical devices designed for single use must not be used multiple times – always follow manufacturer’s instructions.

Cleaning

Cleaning is an important part of stopping the spread of infection and depends on the objects involved and risk of contamination.

When cleaning:

- Most surfaces can be adequately cleaned with warm water and detergent as per manufacturer’s instructions.
- Allow cleaned surfaces to dry completely.
- Detergent solution followed by disinfectant may be appropriate when an infection is known or suspected.

Managing spills

Prompt clean-up of spills (e.g., vomit or diarrhoea) helps to stop infectious agents spreading from the environment to people.

When managing spills:

- Select the appropriate PPE (e.g., gloves and other equipment) depending on the size of the spill
- Immediately wipe up spots and spills smaller than 10cm or cover larger spills with absorbent material
- Dispose of contaminated cleaning materials
- Clean with detergent solution and consider following with disinfectant for infectious or larger spills
- Always perform hand hygiene

Food handling and preparation

Safe food handling is important to prevent food-borne illness.

When handling food:

- Employee must advise their supervisor if suffering from diarrhoea, vomiting, fever, sore throat with fever or jaundice, seek medical advice and not return to work until free of symptoms for 48 hours.
- Employee must advise their supervisor if they have infected skin lesions (e.g., infected skin sore, boil, abrasion, or any discharges from the ears, nose, or eyes) and seek medical advice.
- Employee must advise their supervisor if they believe any food is unsafe to eat.
- Perform hand hygiene before handling food or putting on gloves.
- Perform hand hygiene after using the toilet, smoking, coughing, sneezing, blowing nose, touching face, nose, ears or mouth, handling rubbish or after cleaning.
- Avoid unnecessary contact with ready to eat foods.
- Tie back long hair and cover hair if necessary.
- Secure hair clips, hair pins, buttons on clothes, jewellery, and bandages.
- Ensure bandages or dressings on any exposed parts of the body are covered with a waterproof covering.
- Do not sneeze, blow, cough over unprotected food or surfaces likely to come in contact with food.
- Do not spit, smoke, or use tobacco or similar preparations in area where food is handled.
- Do not touch food after touching earrings, body parts (hair, nose, ear, eye), skin lesions, saliva, mucus, sweat, blood, or money without first performing hand hygiene.
- Do not wear jewellery or lanyards that may come into contact with food.

When preparing food:

- Keep hot food hot and cold food cold